

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

- ⁹ Paris, C. R., Acad. Sci., 163, 1914, (428).
- ¹⁰ Ann. Physik., Leipzig, 29, 1909, (833).
- ¹¹ Astroph. J., Chicago, Ill., 39, 1914, (189).

THE EFFECT OF OXYGEN TENSION ON THE METABOLISM OF CASSIOPEA

By J. F. McClendon

DEPARTMENT OF PHYSIOLOGY, UNIVERSITY OF MINNESOTA AND TORTUGAS LABORATORY,
CARNEGIE INSTITUTION OF WASHINGTON
Communicated by A. G. Mayer, September 21, 1917

It was shown by Verzár (J. Physiol., 45, 1912, 39) that decreased oxygen tension in the blood capillaries decreased the metabolism of muscle but not of salivary glands. The animal died before the oxygen tension in the salivary glands was reduced sufficiently to cause a noticeable fall in metabolism. In order to avoid complications in circulation. as in Verzár's experiments, I used the umbrella of Cassiopea in such a manner as to maintain a thin layer of cells of uniform activity constantly bathed with sea water at 30°. It had been determined in preliminary experiments that a rise of 10° in temperature doubled the metabolism, but that the hydrogen ion concentration could be changed within certain limits without changing the metabolism to a degree that could be measured with certainty. The average of a large number of determinations placed the respiratory quotient at 0.95, but whether it was constant could not be determined. From the heat produced and nitrogen lost it was concluded that proteins with a small admixture of carbohydrates and fats were burned.

Since the temperature was constant, the oxygen-tension was proportional to the O₂-concentration, as determined by the Winkler method, the mean of the values at the beginning and end of the experiment being used. The metabolism was measured by the oxygen used, as that was determined more accurately than the heat and CO₂ eliminated. The metabolism varied with oxygen concentration. This may be true of the cells of all animals. It seems possible that Verzár did not succeed in markedly changing the O₂-tension in the salivary gland, owing to the great store of oxygen in the hemoglobin. There is, however, a distinction between the metabolism of vertebrate muscle cells and Cassiopea. If vertebrates are asphyxiated, the muscles give out lactic acid. A Cassiopea may remain without oxygen for seven hours without giving out CO₂ or any other acid causing a noticeable change in hydrogen ion concentration, although in the presence of O₂ such a change appears in

a few minutes due to elimination of CO_2 . After seven hours without O_2 , nerve conduction and noticeable contractility of muscle returns in thirty seconds after suspending the Cassiopea in air.

Since O₂-tension may affect metabolism, it seems probable that changes in the threshold of stimulation of the respiratory and vasomotor centers may affect metabolism in man and mammals. The details of the experiments will be published elsewhere.